

Distribution of polarization and intensity of radiation across the stellar disk and numerical values of atmospheric characteristics governing this distribution

Bochkarev N., Karitskaya E., Sakhibullin N.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Computations of polarization and intensity of radiation from a unit stellar surface area are presented, as well as a study of the numerical characteristics of atmospheres - single-scattering albedo Ω_λ and the initial source function $\lambda(\delta\lambda)$, which define the polarization behaviour of atmospheres. The radiatively stable models of stellar atmospheres presented by Kurucz et al. (1974) and Kurucz (1979) have been used for calculations. Since the Ω_λ versus optical depth τ_λ dependence is rather weak, it has been assumed that $\Omega_\lambda(\tau_\lambda = \text{const.})$. With a fixed effective temperature T_{eff} maximum values of Ω are characteristic of stars featuring the lowest surface gravity acceleration g . Among stars with radiatively stable atmospheres, maximum values of Ω ($\lambda = 5000 \text{ \AA}$) ≈ 0.4 - 0.6 are exhibited by supergiants with $T_{\text{eff}} = 8000$ - $20\,000 \text{ K}$. The plot of $\Omega(\lambda)$ is characterized by discontinuities at the boundaries of spectral series for hydrogen and, sometimes, for helium. Maximum Ω_λ are attained in the Lyman region of $\lambda = 912$ - 1200 \AA , where Ω_λ can reach the value 0.7 - 0.9 for supergiants, this value being ≥ 0.3 for Main-Sequence stars. For stars with $T_{\text{eff}} \geq 35\,000 \text{ K}$, high values of Ω_λ also are attained for $\lambda < 912 \text{ \AA}$. Within the infrared region, Ω_λ is always small because of bremsstrahlung absorption. A rapid growth of the source function B_λ with $<\lambda$ typical for ultraviolet range (within the Wien part of spectrum), together with high values of Ω_λ results in the strong polarization of emission from a unit stellar surface element, sometimes exceeding the values for the case of a pure electron scattering. For longer wavelengths, where the limb-darkening coefficient is smaller, the plane of polarization abruptly turns 90° in the central parts of the visible stellar disk. © 1985 D. Reidel Publishing Company.

<http://dx.doi.org/10.1007/BF00650117>
